

Recent studies using FerriScan for iron quantification

Inati A et al. [Non-transferrin bound iron in transfused patients with sickle cell disease.](#) *Int J Lab Hematol.* 2010. Feb 25. Epub ahead of print.

Majhail NS et al. [A prospective study of iron-overload management in allogeneic hematopoietic-cell transplant survivors.](#) *Biol Blood Marrow Transplant.* 2010; 16(5):832-7.

Pathare A, et al. [Deferasirox \(Exjade®\) significantly improves cardiac T2* in heavily iron-overloaded patients with beta-thalassemia major.](#) *Ann Hematol* 2010; 89(4):405-9.

Taher A, et al. [Levels of non-transferrin-bound iron as an index of iron overload in patients with thalassaemia intermedia.](#) *Br J Haematol* 2009;146(5):569-572. Jul 13. epub ahead of print.

Taher A, et al. [Magnetic resonance evaluation of hepatic and myocardial iron deposition in transfusion-independent thalassemia intermedia compared to regularly transfused thalassemia major patients.](#) *Am J Hematol.* 2010; 85(4): 288-90.

Patton N, et al. [Observational study of iron overload as assessed by magnetic resonance imaging \(MRI\) in an adult population of transfusion dependent patients with beta thalassaemia: Significant association between low cardiac T2* < 10 ms and the occurrence of cardiac events.](#) *Intern Med J* 2009. May 8. epub ahead of print

Inati A et al. [Absence of cardiac siderosis by MRI T2* despite transfusion burden, hepatic and serum iron overload in Lebanese patients with sickle cell disease.](#) *Eur J Haematol* 2009; 83(6):565-71.

Majhail NS, et al. [Iron-overload after autologous hematopoietic cell transplantation.](#) *Leuk Res* 2009;33(4):578-579.

Olynyk J, et al. [Predicting iron overload in hyperferritinemia.](#) *Clinical Gastroenterology and Hepatology.* 2009;7(3):359-362.